

CONSENT FOR RELEASE OF PENILE SCROTAL WEB (REDUCTION)

Removal of penile scrotal web is usually performed in the operating room under sterile conditions with local anesthesia or conscious sedation. Although strict precautions are taken to prevent infections, no guarantee can be made that this will not occur. Spotting of blood on the dressing may be seen for a few hours following this procedure. Although remote, complications that could occur include excessive bleeding, wound separation, and some degree of localized numbness.

I understand patient cleanliness, dressing changes, applications of wraps, avoidance of trauma, and antibiotics if recommended are essential to a good cosmetic result.

No penetrating sex for 6 weeks, avoid strenuous lifting, no smoking, bed rest for the first 24 hours.

I have read and understand the above, and I hereby consent to this procedure performed by Dr. Harold Reed and his assistant (s) at the Reed Centre for Ambulatory Surgery.

I do not wish to have the tissue analyzed by a Pathologist. Initial: \_\_\_\_\_

I request analysis by a Pathologist and understand my insurance carrier will be billed, if not applicable, I accept the responsibility for any and all lab fees. Initial: \_\_\_\_\_

PATIENT SIGNATURE: \_\_\_\_\_ WITNESS: \_\_\_\_\_

DATE: \_\_\_\_\_ TIME: \_\_\_\_\_